

Original Paper

Efficacy of a Theory-Based Cognitive Behavioral Technique App-Based Intervention for Patients With Insomnia: Randomized Controlled Trial

Nilofar Rajabi Majd¹, MSc; Anders Broström^{2,3}, PhD; Martin Ulander³, PhD; Chung-Ying Lin⁴, PhD; Mark D Griffiths⁵, PhD; Vida Imani⁶, MD; Daniel Kwasi Ahorsu⁴, MPhil; Maurice M Ohayon⁷, PhD; Amir H Pakpour^{1,2}, PhD

¹Social Determinants of Health Research Center, Research Institute for Prevention of Non-Communicable Diseases, Qazvin University of Medical Sciences, Qazvin, Iran

²Department of Nursing, School of Health and Welfare, Jönköping University, Jönköping, Sweden

³Department of Clinical Neurophysiology, Linköping University Hospital, Linköping, Sweden

⁴Department of Rehabilitation Sciences, Hong Kong Polytechnic University, Hung Hom, China (Hong Kong)

⁵International Gaming Research Unit, Psychology Department, Nottingham Trent University, Nottingham, United Kingdom

⁶Pediatric Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁷Division of Public Mental Health and Population Sciences, Stanford Sleep Epidemiology Research Center, Stanford University, Palo Alto, CA, United States

Corresponding Author:

Chung-Ying Lin, PhD

Department of Rehabilitation Sciences

Hong Kong Polytechnic University

ST534 Department of Rehabilitation Sciences

Hung Hom,

China (Hong Kong)

Phone: 852 27666755

Email: cylin36933@gmail.com

Abstract

Background: Sleep hygiene is important for maintaining good sleep and reducing insomnia.

Objective: This study examined the long-term efficacy of a theory-based app (including cognitive behavioral therapy [CBT], theory of planned behavior [TPB], health action process approach [HAPA], and control theory [CT]) on sleep hygiene among insomnia patients.

Methods: The study was a 2-arm single-blind parallel-group randomized controlled trial (RCT). Insomnia patients were randomly assigned to a treatment group that used an app for 6 weeks (ie, CBT for insomnia [CBT-I], n=156) or a control group that received only patient education (PE, n=156) through the app. Outcomes were assessed at baseline and 1 month, 3 months, and 6 months postintervention. Primary outcomes were sleep hygiene, insomnia, and sleep quality. Secondary outcomes included attitudes toward sleep hygiene behavior, perceived behavioral control, behavioral intention, action and coping planning, self-monitoring, behavioral automaticity, and anxiety and depression. Linear mixed models were used to evaluate the magnitude of changes in outcomes between the two groups and across time.

Results: Sleep hygiene was improved in the CBT-I group compared with the PE group ($P=.02$ at 1 month, $P=.04$ at 3 months, and $P=.02$ at 6 months) as were sleep quality and severity of insomnia. Mediation analyses suggested that perceived behavioral control on sleep hygiene as specified by TPB along with self-regulatory processes from HAPA and CT mediated the effect of the intervention on outcomes.

Conclusions: Health care providers might consider using a CBT-I app to improve sleep among insomnia patients.

Trial Registration: ClinicalTrials.gov NCT03605732; <https://clinicaltrials.gov/ct2/show/NCT03605732>

(*J Med Internet Res* 2020;22(4):e15841) doi: [10.2196/15841](https://doi.org/10.2196/15841)

KEYWORDS

app-based intervention; cognitive behavioral therapy; insomnia; sleep hygiene; theory of planned behavior